Fertility/IVF referral pathway involving primary care (part 1)

Female

investigations

also required

KEY

Primary Care

Secondary Care

Oxfordshire
Clinical
Commissioning
Group

Private Care

Primary Care investigation of all couples **STARTS** with

Couple sees GP with failure to conceive

See **NICE guidelines**

Male investigations

 Analysis of seminal fluid. This should be the first step in the investigation of infertility since in the absence of sperm the fertility appointment is not required.

Method

- 3 days abstinence from intercourse
- collection of specimen of semen in a sterile plastic container after masturbation
- examine within time set by clinic prior appointment

See this link for options and forms

Referral made to

IVF UNIT of CHOICE (see part 2)

Please use OCCG guidelines for eligibility

criteria for NHS funded treatment and the

referral form

Options available on the CCG website here

Female investigations

- Assessment of ovulation mid-luteal progesterone taken seven days prior to menses e.g. day 21 of 28 day cycle or day 25 of 32 day cycle. For **amenorrhoea** just state this at the time of the tests.
- Other hormonal tests LH, testosterone,
- FSH, Prolactin performed on day 2 5 of cycle. See above for amenorrhoea.
- Rubella status (if not already known)
- In date cervical smear
- Chlamydia swab
- Pelvic/TV USS if available

Refer all couples to

NHS Fertility Clinic

(Via C&B) with above

test results

NO SPERM

See this link

Or previously proven inability to conceive

Azoospermatic couples who meet the funding criteria go straight to IVF with the female partner's tests, which are required prior to referral.

Normal/Low sperm count See this link

- Normal / low sperm count
- Abnormal bloods and/or
- Abnormal scan and/or
- No conception after one year of regular SI (but see previous advice)

Patients seen and treated in

NHS FERTILITY CLINIC

(see part 2)

CONCEPTION and/or END OF TREATMENT

Approved CRG July 2015 Review July 2018 Dr S Hayles