

## OCCG SERVICE SPECIFICATION (2021/22)

### Primary Care Service for Skin Cancers: Dermatology Shared Care Monitoring for Melanoma, Lichen Sclerosus and Squamous Cell Carcinoma

#### 1. Background

For patients who have had a diagnosis of Melanoma, Squamous Cell Carcinoma, Vulval Lichen Sclerosus or Vulval Lichen Planus, in-line with national guidelines, it is important that there is ongoing monitoring to detect any potential malignant change/recurrence.

For some of this activity it is appropriate that the care is undertaken in primary care under shared care arrangements. A copy of the shared care protocol (see from appendix 2 onwards) will be sent to the practice/GP when a patient is started on this pathway by the dermatology department. Most will eventually be discharged back to primary care as the majority of care for patients with a diagnosis of vulval lichen sclerosus or vulval lichen planus is provided through the GP, also in line with national guidance.

Since 2009/10 the dermatology department has been reviewing all patients with a diagnosis of Lichen Sclerosus. A large number have been discharged back to primary care, as they have been assessed as being at very low risk of complications. Patients are advised to visit their GP annually for review and to report any skin changes to their GP. These patients are not covered by shared care as they are discharged back to primary care.

In patients identified by the hospital consultant as presenting with symptoms that are difficult to control or have poor response to treatment or concerns about progressive severe scarring, the management can be undertaken through a shared care arrangement. For such patients, practices are eligible for one annual payment under service, for two reviews per annum.

Patients diagnosed with Stage IB and IIA **Malignant Melanoma** (Level III, Grade B) can be managed through shared care arrangements following diagnosis. Patients will be seen twice a year in primary care and twice a year in secondary care for the first three years. In the fourth and fifth year the patient will be seen once in primary care and once by a secondary care consultant.

A new low risk group for **Malignant Melanoma** (Stage IA Melanoma) has been identified which will only need following up at 4 monthly intervals for one year, having two visits to the hospital and one to the GP before being discharged from formal follow up in secondary care.

Shared care arrangements for medium risk **Squamous Cell Carcinomas** are being introduced in line with the national guidelines which specify 6 monthly attendances with the GP over two years. During that time, they will also be seen by the hospital 6 monthly.

No part of this specification by commission, omission or implication defines or redefines essential or additional services.

#### 2. Service Scope

See appendices for guidelines for Lichen Sclerosus, Lichen Planus, Squamous Cell Carcinoma and Melanoma.

#### 3. Patient Re-call

Practices are required to create and maintain a register of patients treated under the shared care protocol and run regular searches identifying patients due for review. The searches used by the CSU for the extraction of data would help to identify the search population – please contact [scwcsu.primarycareanalytics@nhs.net](mailto:scwcsu.primarycareanalytics@nhs.net) for how to access.

Patients should be invited in writing to attend the practice for their review in line with the shared care protocol. The letter should detail the purpose of the visit and recommend attendance.

#### 4. Coding

Please see Appendix 1 for the codes to be used.

The coding required for payment includes three codes:

1. The shared care code
2. A diagnosis code
3. Dermatology examination code

#### 5. Payment

Practices will receive payments as indicated in the table below for reviews undertaken under the shared care arrangements.

Please note that the frequency of the review is dependent upon the diagnosis:

Diagnosis	Frequency of Review	Payment per review
Lichen Sclerosus	Two reviews per annum in primary care	£56.10
Erosive Lichen Planus	Two reviews per annum in primary care	£56.10
Malignant Melanoma – Stage IA	One review in primary care Duration 1 year	£56.10
Malignant Melanoma – Stage IB and IIA	Two reviews in primary care per annum for 3 years / one review in primary care in years 4 and 5	£56.10
Squamous Cell Carcinoma	Two reviews in primary care per annum. Duration 2 years	£56.10

Payment will be made quarterly based on actual activity carried out as reflected in quarterly activity monitoring reports to the CCG as per clause 6 below.

#### 6. Monitoring

Data will be extracted from the Emis practice system by SCWCSU around the 15th of the month following the end of each quarter to monitor activity. Vision practices are requested to submit the necessary evidence for payment in the form of the search output from the clinical system and submit to the SCWCSU by 15th of the month following the end of each quarter during the year. (Deadlines to be confirmed each quarter by SCWCSU).

#### 7. Termination

This service will terminate on 31<sup>st</sup> March 2022. For termination ahead of this 3 month written notice must be given.

#### 8. Contact

Contact for queries: [paul.kettle1@nhs.net](mailto:paul.kettle1@nhs.net)  
 Clinical lead: [shelley.hayles@nhs.net](mailto:shelley.hayles@nhs.net)

**APPENDIX 1**

**Oxfordshire CCG**

Data Collection Specification for Primary Care for Skin Cancers: Dermatology Shared Care Monitoring for Melanoma, Lichen Sclerosus and Squamous Cell Carcinoma 2021/22

*Lichen Planus*

	<b>Item</b>	<b>Read Code and Description</b>	<b>SNOMED Codes</b>	<b>Notes</b>
Search Population	Vulval Lichen Planus and shared care arrangement with the specialist	M1702 Lichen planus atrophicus 66S2. Shared care - specialist / GP	25858008 268529002	Any record of both codes
For payment	Dermatology exam	2F... O/E - dermatology exam.	271898004	Maximum 2 claims in the year

*Lichen Sclerosus*

	<b>Item</b>	<b>Read Code and Description</b>	<b>SNOMED Codes</b>	<b>Notes</b>
Search Population	Vulval Lichen Sclerosus and shared care arrangement with the specialist	M2102 Lichen sclerosus et atrophicus 66S2. Shared care - specialist / GP	25674000 268529002	Any record of both codes
For payment	Dermatology exam	2F... O/E - dermatology exam.	271898004	Maximum 2 claims in the year

*Malignant Melanoma*

	<b>Item</b>	<b>Read Code and Description</b>	<b>SNOMED Codes</b>	<b>Notes</b>
Search Population	Malignant Melanoma in the last 5 years and shared care arrangement with the specialist in the last year	B32% Malignant melanoma of skin 66S2. Shared care - specialist / GP	93655004 268529002	Latest MM code which will be assigned to Staging categories below
	Stage IA	B328. Malignant melanoma stage IA	956331000000107	1 Review in Primary Care
For payment	Dermatology exam	2F... O/E - dermatology exam.	271898004	
	Stage IB or IIA	B329. Malignant melanoma stage IB	956351000000100 956371000000109	Years 1-3: 2 reviews

		B32A. Malignant melanoma stage IIA		Years 4 & 5: 1 review
For payment	Dermatology exam		271898004	

*Squamous Cell Carcinoma*

	Item	Read Code and Description	SNOMED Codes	Notes
Search population	Squamous Cell Carcinoma in the last 2 years and shared care arrangement with the specialist in the last year also in the last 2 years	B338. Squamous cell carcinoma of skin 66S2. Shared care - specialist / GP	254651007 268529002	Maximum 2 claims in the year
For payment	Dermatology exam	2F... O/E - dermatology exam.	271898004	

Data extraction is cumulative throughout the fiscal year taken at the end of each quarterly period. In all cases, patients who have died or left during the reporting period, but who have received a service within the practice will be included.

Temporary residents will not be included.

*Please note SNOMED Codes are currently indicative, subject to data quality checks.*

## APPENDIX 2

### VULVAL LICHEN SCLEROSUS: PRIMARY CARE MANAGEMENT

The dermatology dept. has been reviewing the follow up of Vulval Lichen Sclerosus patients in view of recent national guidelines in patients diagnosed with Lichen Sclerosus and concluded that many lower risk women would be adequately managed by annual review in primary care.

Patients whose disease is stable and low risk will be discharged from the clinic after seeing the consultant to discuss this. They will be advised to self-examine and visit their GP if they develop persistent ulcers or nodules that don't respond to three weeks twice daily topical treatment with Dermovate.

All patients who have used topical steroid during the previous year are recommended to be reviewed annually and examined in primary care. *Please see below for more detail.*

Patients should be referred back urgently into the specialist service if there are concerns.

Below is information for GPs

  
Department of Dermatology  
The Churchill Hospital  
Tel: 01865 228266  
Fax: 01865 228260

### VULVAL LICHEN SCLEROSUS: COMMUNITY CARE

#### Lichen Sclerosus

Your patient has been attending our clinic with Lichen Sclerosus of the vulva and/or perianal area. This condition is a chronic one and is associated with a 3-5% risk of malignant change which may occur as early as the 30s. We are currently recommending that following treatment, patients are followed up yearly for signs of early malignant change. We are also encouraging women to self-examine and report any changes.

Community care is suitable for women with well controlled Lichen Sclerosus (requiring less than 30g Dermovate ointment or equivalent in any 6 month period), those who find travel to a hospital clinic too difficult or for those who express a preference for GP care. We propose that follow-up care should be carried out by the General Practitioner at yearly intervals. In case your experience with this disease is limited we have produced this advice sheet to help you. *Please contact us or refer back if you have any concerns.*

#### ***Vulval appearances and symptoms to expect in Lichen Sclerosus:***

- The disease can present either as a localised patchy problem, or involvement can be extensive affecting the entire vulva/perineum, typically extending to the perianal area
- Plaques are usually white and generally thin and atrophic (like cigarette paper)
- Purpura and haemorrhage are common features
- Architectural change is common and there may be labial fusion with a contracted introitus, and the clitoris may be buried
- Fissures are common, but must be seen to heal
- Secondary infection with candida (which may be clinically atypical) and bacteria may cause worsening of symptoms. Please do a **vulval** swab and treat as appropriate if you suspect this
- Remember that many patients are post-menopausal and may also need local oestrogen

**What to look for:**

- Erosions which do not respond to topical application of Dermovate Ointment twice daily for three weeks should be referred urgently for review at the vulval clinic.
- Hyperkeratotic areas or fissured areas that do not respond to Dermovate Ointment twice daily for three weeks need an urgent referral
- Nodule formation is a very suspicious sign and needs urgent referral. *If a tumour is strongly suspected urgent referral via the 2 week wait to gynaecology is preferable to avoid delay in treatment.*
- Lichen Sclerosus has a strong clinical association with other autoimmune diseases such as thyroid disease, pernicious anaemia, vitiligo and diabetes. Please remain vigilant that these can arise at any stage and may require treatment and monitoring.

**Dr Susan Cooper**  
**Consultant Dermatologist**

**Sister Susan Booker**  
**Nurse Practitioner**

SMC March 2011

-----  
**VULVAL LICHEN SCLEROSUS: SHARED CARE**

Below is the information for GPs:

  
Department of Dermatology  
The Churchill Hospital  
Tel: 01865 228266  
Fax: 01865 228260

## **VULVAL LICHEN SCLEROSUS: SHARED CARE**

### **Lichen Sclerosus**

Your patient has been attending our clinic with Lichen Sclerosus of the vulva and/or perianal area. This condition is a chronic one and is associated with a 3-5% risk of malignant change which may occur as early as the 30s.

We propose that shared follow-up care should be carried out between the hospital and the General Practitioner and should be on the basis of hospital review every eighteen months with the rest of the six monthly reviews be carried out in the community. When the disease is in clinical remission (symptoms well controlled and requiring less than 30g Dermovate ointment in any 6 month period) we will recommend that all reviews are done in the community. Community care may also be more suitable for those who find travel to hospital difficult or express a preference for community care. Patients can be referred back urgently if there are any concerns.

In case your experience with this disease is limited we have produced this advice sheet to help you.

### ***Vulval appearances and symptoms to expect in Lichen Sclerosus:***

- The disease can present either as a localised patchy problem, or involvement can be extensive affecting the entire vulva/perineum, typically extending to the perianal area
- Plaques are usually white and generally thin and atrophic (like cigarette paper)
- Purpura and haemorrhage are common features
- Architectural change is common and there may be labial fusion with a contracted introitus, and the clitoris may be buried
- Fissures are common, but must be seen to heal
- Secondary infection with candida (which may be clinically atypical) and bacteria may cause worsening of symptoms. Please do a **vulval** swab and treat as appropriate if you suspect this

- Remember that many patients are post-menopausal and may also need local Oestrogen

***What to look for:***

- Erosions which do not respond to topical application of Dermovate Ointment twice daily for three weeks should be referred urgently for review at the vulval clinic.
- Hyperkeratotic areas or fissured areas that do not respond to Dermovate Ointment twice daily for three weeks need an urgent referral
- Nodule formation is a very suspicious sign and needs urgent referral. If a tumour is strongly suspected referral via the 2 week wait to gynaecology is more appropriate to avoid delays in treatment.
- Lichen Sclerosus has a strong clinical association with other autoimmune diseases such as thyroid disease, pernicious anaemia, vitiligo and diabetes. Please remain vigilant that these can arise at any stage and may require treatment and monitoring.

**Dr Susan Cooper**  
**Consultant Dermatologist**

**Sister Susan Booker**  
**Nurse Practitioner**

SMC March 2011

~~~~~



## APPENDIX 3

### VULVAL LICHEN PLANUS – SHARED CARE

Some **erosive Lichen Planus** may be suitable for shared care, being seen at 6 monthly intervals with every second or third visit to a consultant.

$\alpha\beta\gamma$

Department of Dermatology  
The Churchill Hospital  
Tel: 01865 228266  
Fax: 01865 228260

## VULVAL LICHEN PLANUS: SHARED CARE

### Lichen Planus

Your patient has been attending our clinic with Lichen Planus of the vulva and/or perianal area. She is currently in clinical remission; however this condition is a chronic one and is associated with a 3-5% lifetime risk of malignant change which may occur as early as the 30s.

We propose that shared follow-up care should be carried out between the hospital and the General Practitioner and should be on the basis of hospital review every eighteen months with the rest of the six monthly reviews be carried out in the community. When your patient is in clinical remission requiring minimal treatment **we will recommend that all follow up takes place in the primary care**. In case your experience with this disease is limited we have produced this advice sheet to help you. *Please contact us or refer back if you have any concerns.*

### ***Vulval appearances and symptoms to expect in Lichen Planus:***

- The disease usually presents with erosions at the vaginal introitus that are typically edged by a lacy-white border. These erosions often persist.
- In some cases (hypertrophic disease) thickened white plaques may develop
- Architectural change is common and there may be labial fusion with a contracted introitus, and the clitoris may be buried.
- Secondary infection with candida (which may be clinically atypical) and bacteria may cause worsening of symptoms. Please do a **vulval** swab and treat as appropriate if you suspect this.
- Remember that many patients are post-menopausal and may also need local Oestrogen

### ***What to look for:***

- New hyperkeratotic areas that do not respond to Dermovate ointment twice daily for three weeks.
- Very persistent, thickened or enlarging erosions.
- Nodule formation is a very suspicious sign and needs urgent referral. *If a tumour is strongly suspected then referral via the 2 week wait to gynaecology is more appropriate to avoid delays in treatment.*
- Rarely the narrowing of the vaginal introitus may be so severe that urinary retention may occur.
- Lichen Planus can occur at other body sites eg the oral mucosa, scalp, extragenital skin and nails.
- Finally Lichen Planus has a strong clinical association with other autoimmune diseases such as thyroid disease, pernicious anaemia, vitiligo and diabetes. Please remain vigilant that these can arise at any stage and may require treatment and monitoring.

**Dr Susan Cooper**  
**Consultant Dermatologist**

**Sister Susan Booker**  
**Nurse Practitioner**



## VULVAL LICHEN PLANUS – PRIMARY CARE MANAGEMENT

 $\alpha\beta\gamma$ 

Department of Dermatology

The Churchill Hospital

Tel: 01865 228266

Fax: 01865 228260

**VULVAL LICHEN PLANUS: COMMUNITY CARE****Lichen Planus**

Your patient has been attending our clinic with Lichen Planus of the vulva and/or perineum. She is currently in clinical remission, however this condition is a chronic one and is associated with an estimated 3-5% risk of malignant change which may occur as early as the 30s. We are currently recommending that following treatment, patients are followed up at yearly intervals for life for signs of early malignant change.

Community care is suitable for women with well controlled Lichen Planus (typically requiring less than 30g Dermovate ointment or equivalent in any 6 month period), those who find travel to a hospital clinic difficult or for those who express a preference for GP care. We propose that follow-up care should be carried out by the General Practitioner at six monthly intervals. In case your experience with this disease is limited we have produced this advice sheet to help you. *Please contact us or refer back if you have any concerns.*

***Vulval appearances and symptoms to expect in Lichen Planus:***

- The disease usually presents with erosions at the vaginal introitus that are typically edged by a lacy-white border. These erosions often persist.
- In some cases (hypertrophic disease a) thickened white plaques may develop
- Architectural change is common and there may be labial fusion with a contracted introitus, and the clitoris may be buried.
- Secondary infection with candida (which may be clinically atypical) and bacteria may cause worsening of symptoms. Please do a **vulval** swab and treat as appropriate if you suspect this.
- Remember that many patients are post-menopausal and may also need local Oestrogen

***What to look for:***

- New hyperkeratotic areas that do not respond to Dermovate ointment twice daily for three weeks.
- Very persistent, thickened or enlarging erosions.
- Nodule formation is a very suspicious sign and needs urgent referral. *If a tumour is strongly suspected urgent referral via the 2 week wait to gynaecology is preferable to avoid delay in treatment.*
- Rarely the narrowing of the vaginal introitus may be so severe that urinary retention may occur.
- Lichen Planus can occur at other body sites e.g. the oral mucosa, scalp, extragenital skin and nails.
- Lichen Planus has a strong clinical association with other autoimmune diseases such as thyroid disease, pernicious anaemia, vitiligo and diabetes. Please remain vigilant that these can arise at any stage and may require treatment and monitoring.

**Dr Susan Cooper**  
**Consultant Dermatologist**

**APPENDIX 4**

Melanoma Follow Up

Stage IA Melanoma – Low Risk

Under this service practices will undertake one annual review in the 12 months following diagnosis of a low risk Melanoma for those patients that are being treated under a shared care protocol as below:

Melanoma Follow Up

Stage IB and IIA Melanoma

The patient will be seen in the outpatient clinic initially then further appointments will usually alternate between the Specialist and GP according to the stage of disease at diagnosis. The patient should check the original Melanoma site once a week for the first year then once a month after that. The frequency of the hospital and GP visits depends on the thickness of the Melanoma. The scar, surrounding skin, regional draining lymph nodes and general skin should be examined. Details of the likely frequency of visits is detailed below:

|                                                                                            | Stage IA Melanoma                                                                                                                                                                  | Stage IB and IIA Melanoma                                                                                                                                                                                                                                                                                                                 | Stage IIB, IIC and IIIA Melanoma                                                                                                                | Stage IIIB, IIIC and IV Melanoma                                                                                                                                                                        |
|--------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                                            | Level III, Grade B                                                                                                                                                                 | Level III, Grade B                                                                                                                                                                                                                                                                                                                        | Level III, Grade B                                                                                                                              | Level III, Grade B                                                                                                                                                                                      |
|                                                                                            | <b>Shared Care Potential</b>                                                                                                                                                       | <b>Shared Care Potential</b>                                                                                                                                                                                                                                                                                                              | Secondary Care Led                                                                                                                              | Secondary Care Led                                                                                                                                                                                      |
| Recommendation Ref: BAD Guidelines for mgt of cutaneous Melanoma 2010, J.R. Marsden et al. | Three visits over 12 months (one in <b>primary care</b> and two in secondary care) are suggested to teach self-examination, and then they may be discharged from regular follow up | Patients should learn how to self-examine for locoregional metastasis and new primaries, and understand how to access the follow-up team promptly for suspect recurrence, They should be seen every 3 months for 3 years, then 6 monthly to 5 years alternating between primary & secondary care. No routine investigations are required. | Patients should be taught self-examination and be seen 3 monthly for 3 years, and 6 monthly to 5 years. No routine investigations are required. | Many patients will be eligible for adjuvant trials. Those outside of trials should be seen 3 monthly for 3 years from the date of staging, 6 monthly to 5 years, then annually to 10 years by an SSMdT. |

Patients will, as far as possible, be instructed in self-examination and receive written information about Melanoma, the contact details of the skin cancer nurse specialist & clear instructions and actions to take should they suspect new or recurrent disease. All patients are at risk of further skin cancers & pre-cancerous skin disease such as actinic (solar) keratosis & Bowen’s disease. They will receive written information on sun avoidance measures.

The original site i.e. scars and surrounding skin as well as the regional draining lymph nodes should be examined. The patient should be asked about and examined for other suspicious skin lesions. All patients suitable for shared care follow-up will be given a patient-held follow-up record documenting their details including diagnosis, specialist contact details, follow-up regimen & chart to record clinical findings. The patient is instructed to bring this to each GP & specialist follow-up appointment.

~~~~~

## **Appendix 5**

### **Squamous Cell Carcinoma (SCC)**

SCC shared care guidance has recently been agreed nationally, and some patients with SCCs can be seen in primary care under this. This will now be paid for as part of this service in a similar way to the malignant Melanoma payment.

Early detection and treatment improves survival of patients with recurrent disease. Seventy-five percent of local recurrences and metastases are detected in 2 years and 95% are detected within 5 years. The Department of Dermatology or Plastic Surgery will stratify SCC patients into 3 risk categories: low, medium & high based on the anatomical site affected, the size of the tumour, its underlying aetiology, degree of histological differentiation and host immunosuppression. The risk of metastasis is related to its site - in order of increasing metastatic potential:

1. SCC arising at sun-exposed sites excluding lip and ear.
2. SCC of the lip.
3. SCC of the ear.
4. Tumours arising in non-sun-exposed sites (e.g. perineum, sacrum, sole of foot).
5. SCC arising in areas of radiation or thermal injury, chronic draining sinuses, chronic ulcers, chronic inflammation or Bowen's disease.

Patients will, as far as possible, be instructed in self-examination and receive written information about SCC, the contact details of the skin cancer nurse specialist & clear instructions and actions to take should they suspect new or recurrent disease. All patients are at risk of further skin cancers & pre-cancerous skin disease such as actinic (solar) keratoses & Bowen's disease. They will receive written information on sun avoidance measures.

#### **Low risk**

SCCs judged to be of low risk of local recurrence and/or metastasis will be reviewed in secondary care, usually the Department of Dermatology, 3 months after their definitive treatment and then discharged. They do not need regular follow up thereafter.

Low risk SCCs are defined as those arising in patients with no background immune dysfunction on sun-exposed sites, except the lip & ear, measuring less than 20mm in diameter or 4mm in thickness that are well differentiated

#### **Medium risk**

Those judged to be of medium risk will be followed up for 2 years at 3 monthly intervals i.e. twice a year by the GP and twice a year by the hospital on an alternating basis. The original site i.e. scars and surrounding skin as well as the regional draining lymph nodes should be examined. The patient should be asked about and examined for other suspicious skin lesions. All patients suitable for shared care follow-up will be given a patient-held follow-up record documenting their details including diagnosis, specialist contact details, follow-up regimen & chart to record clinical findings. The patient is instructed to bring this to each GP & specialist follow-up appointment.

#### **High risk**

These patients will be followed up for 2 years with shared care with primary care every 6 months and secondary care seeing them at the 3 month interim appointment.